

Metric Fuzziness and Eigenvalue Theory

Gunjan Deva Sarma^{1*}, Dr. Manmohan Das²

^{1*}Department of Mathematics, Bhattadev University, Bajali, Assam, India, 781326, Mail- sarmagunjandev@gmail.com

²Department of Mathematics, Bhattadev University, Bajali, Assam, India, 781326, Mail- mdas.bajali@gmail.com

Abstract:

This article is dedicated to the exploration of fuzzy eigenvalues and fuzzy eigenvectors within the context of a fuzzy metric space. To facilitate this discussion, we introduce a specific metric for this space. Furthermore, we provide comprehensive definitions for fuzzy eigenvalues and fuzzy eigenvectors, focusing on their application to fuzzy square matrices. In the course of our exploration, we establish a series of theorems pertaining to fuzzy eigenvalues and eigenvectors within a fuzzy metric space. To enhance understanding, we illustrate these theorems with practical examples.

Keywords: Fuzzy eigenvalue, Fuzzy eigenvector, Fuzzy eigen space, metric space in fuzzy.

1. INTRODUCTION

In 1975, the concept of a fuzzy metric space was initially presented by Kramosil and Michálek in their seminal work [1]. This innovative concept extends beyond the traditional statistical (probabilistic) metric space, offering a more flexible framework for distance measurement. Subsequently, Grabiec's work [2] focused on establishing the notion of completeness within the realm of fuzzy metric spaces, a pivotal property for studying convergence and continuity. This field has since evolved, with researchers exploring two distinct approaches to address the challenges and applications of fuzzy metric spaces. There are two primary approaches in the study of fuzzy metric spaces. One approach, as pioneered by Kaleva in 1984 [6], involves the use of fuzzy numbers to define metrics in conventional spaces. Following Kaleva's work, several researchers have delved into this approach, exploring topics such as fuzzy normed spaces, fuzzy topology derived from fuzzy metric

spaces, fixed point theorems, and other properties of fuzzy metric spaces. Notable contributions include the research of Felbin in 1992 [7], George in 1994 [8], George in 1997 [9], Gregori in 2000 [10], and Hadzic in 2002 [11], among others.

The second approach focuses on utilizing real numbers to quantify the distances between fuzzy sets.

The sources related to this particular approach can be found in works by Dia in 1990 [5], Chaudhuri in 1996 [4], Boxer in 1997 [3], Fan in 1998 [12], Voxmam in 1998 [15], Przemyslaw in 1998 [14], and Brass in 2002 [13]. The findings of these studies have found practical applications in various real-world challenges within fuzzy environments. However, it's worth noting that different problems often require distinct measurement approaches. In other words, there is no universal measure that can be applied across all types of fuzzy environments.

2. PRELIMINARIES Definition:1

If $A \in M_X$ and $x \in \mathbb{R}^n$, then \exists a constant $\lambda \in [0,1]$ such that $Ax \in \lambda x$. Then λ is called an eigenvalue of matrix A , along with its associated eigenvector x .

Definition:2

For $A \in M_X$, $\sigma(A) = \{\lambda : Ax = \lambda x \text{ has a solution for a non-zero vector } x\}$, is referred to as the spectrum of A .

$$\rho(A) = \sup_{\lambda \in \sigma(A)} |\lambda|$$

$\rho(A)$; is called the spectral radius.

The characteristic polynomial:

The polynomial $(A - \lambda I)$ has a non-trivial solution. i.e. $\det(A - \lambda I) = 0$

Thus $\det(A - \lambda I)$ must be a polynomial in λ .

Definition:3

Let D represent the collection of all closed and bounded intervals $X = [a_1, a_2]$ on real numbers. For $\bar{X}, \bar{Y} \in D$, we define $\bar{X} \leq \bar{Y}$ if $p_1 \leq q_1$ and $p_2 \leq q_2$.

$d(\bar{X}, \bar{Y}) = \max(|p_1 - q_1|, |p_2 - q_2|)$, when $\bar{X} = [p_1, p_2]$ and $\bar{Y} = [q_1, q_2]$. It is a well-established fact that the metric space (D, d) is complete.

Definition [16]:4

Assume that X is a nonempty set, and $d_S: PS(X) \times PS(X) \rightarrow S^+(R)$ is a function. $(PS(X), d_S)$ is termed a fuzzy metric space provided that, for every $\{(x_s, \lambda_s), (y_s, \gamma_s), (z_s, \rho_s)\} \subset PS(X)$, d_S meet the following set of three conditions,

- (1) Non-negative: $d_S((x_s, \lambda_s), (y_s, \gamma_s)) = 0$ iff $x_s = y_s$ and $\lambda_s = \gamma_s = 1$;
- (2) Symmetric: $d_S((x_s, \lambda_s), (y_s, \gamma_s)) = d_S((y_s, \gamma_s), (x_s, \lambda_s))$;
- (3) Triangular inequality: $d_S((x_s, \lambda_s), (z_s, \rho_s)) < d_S((x_s, \lambda_s), (y_s, \gamma_s)) + d_S((y_s, \gamma_s), (z_s, \rho_s))$.

d_S is termed as fuzzy metric defined in $PS(X)$ and $d_S((x_s, \lambda_s), (y_s, \gamma_s))$ is referred to as a fuzzy distance between the two fuzzy points.

Definition:5

Let $\bar{A} \in M_X$, then $\rho_{\bar{A}}(\lambda) = |\lambda I - \bar{A}|$ is called the characteristic polynomial of \bar{A} . Its roots are called the latent value of \bar{A} .

Theorem - 1:

Let $\bar{A} \in M_X$. The collection of fuzzy eigenvectors corresponding to a specific fuzzy eigenvalue forms a subspace within the vector space \mathbb{C}_n .

Proof:

Let $\lambda \in \sigma(\bar{A})$ be a fuzzy eigenvalue of a fuzzy square matrix \bar{A} . The set of fuzzy eigenvectors associated with λ is the kernel of the matrix $(\bar{A} - \lambda)$. Once more, the kernel of any fuzzy matrix constitutes a subspace within the fundamental vector space. Hence the result.
(Thus, we get the following definition)

Definition:

Let $\bar{A} \in M_X$ and let $\lambda \in \sigma(\bar{A})$. The kernel of $(\bar{A} - \lambda)$ is referred to as the fuzzy eigenspace of \bar{A} corresponding to λ .

Theorem – 2:

Let $\bar{A} \in M_X$, then fuzzy eigenvectors corresponding to distinct fuzzy eigenvalues exhibit linear independence. Suppose $\lambda \in \sigma(\bar{A})$ with fuzzy eigenvectors x is different from the set of fuzzy eigenvalues $\{\mu_1, \mu_2, \dots, \mu_k\} \in \sigma(\bar{A})$ and V_{μ} is the subspace generated by the associated fuzzy eigenspace. Hence $x \notin V_{\mu}$.

Proof:
Let us consider $\mu, \lambda \in \sigma(\bar{A})$ with $\mu \neq \lambda$ be the fuzzy eigenvalue corresponding to the fuzzy eigen vectors x and y respectively. Then vectors are L.D, i.e. $y = cx$

$$\begin{aligned} \text{Then } \mu y &= \mu cx = c\mu x \\ &= cAx \\ &= A(cx) \\ &= Ay \\ &= \lambda y \end{aligned}$$

This statement presents an inconsistency. And hence (i) is said to be true.

Consider the opposite scenario, where it is assumed that $x \in V_{\mu}$. Thus

$$x = a_1y_1 + \dots + a_my_m;$$

where $\{y_1, \dots, y_m\}$ are L.I in V_{μ} . Every one of the vectors y_i is eigen vector. Let's assume the relevant fuzzy eigenvalues, represented as μ_j .

$$\text{i.e. } Ay_i = \mu_{j_i} y_i, \text{ for each } i = 1, 2, \dots, m$$

$$\begin{aligned} \text{Then } \lambda x &= Ax \\ &= A(a_1y_1 + \dots + a_my_m) \\ &= a_1\mu_{j_1}y_1 + \dots + a_m\mu_{j_m}y_m \end{aligned}$$

If $\lambda \neq 0$, we have

$$x = a_1y_1 + \dots + a_my_m$$

$$= a_1 \mu_{j1} y_1 + \dots + a_m \mu_{jm} y_m$$

We know from the above that at least two of the co-efficient a_i must be non-zero.

Consequently, we encounter two distinct representations of a single vector through linearly independent vectors, which is an infeasible scenario.

On the other hand, if $\lambda = 0$, Then

$$a_1 \mu_{j1} y_1 + \dots + a_m \mu_{jm} y_m = 0$$

Which is also impossible. Thus (ii) is proved.

Illustration-1 of theorem-2:

Let $A = \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \end{bmatrix}$, where $a, b, c, d \in [0,1]$ be a fuzzy square matrix.

Then $\det(\lambda I - A) = \det \begin{pmatrix} \lambda - \bar{a} & -\bar{b} \\ -\bar{c} & \lambda - \bar{d} \end{pmatrix}$
 $= (\lambda - \bar{a})(\lambda - \bar{d}) - \bar{b}\bar{c}$
 $= \lambda^2 - (\bar{a} + \bar{d})\lambda + \bar{d}\bar{b} - \bar{b}\bar{c}$
 \therefore the fuzzy eigen values will be the root of the equation (1).

$$\bar{a} + \bar{d} \pm \sqrt{(\bar{a} + \bar{d})^2 - 4(\bar{d}\bar{b} - \bar{b}\bar{c})}$$

2

$$\bar{a} + \bar{d} \pm \sqrt{(\bar{a} + \bar{d})^2 + 4\bar{b}\bar{c}}$$

Regarding this quadratic, three potential outcomes are under consideration:

- i) A pair of genuine solutions
 - ii) A pair of complex solutions
- Distinct values Identical values

Illustration-2:

Consider a fuzzy matrix denoted as \bar{A} with a rank unity. In such a case, there exist two vectors $w, z \in \mathbb{C}_n$ In this context

$$= wz^T \quad \bar{A}$$

To determine the spectrum of \bar{A} , we examine the fuzzy ch. Polynomial as $Ax = \lambda x$

$$\Rightarrow \langle z, x \rangle w = \lambda x$$

For this, we see that $x = w$ is one fuzzy eigen vector with fuzzy eigen value $\langle z, x \rangle$. If $z \perp x$, x is a fuzzy eigen vector pertaining to the eigen value 0.

$$\therefore \sigma_A = \{ \langle z, x \rangle, 0 \}$$

Hence, the ch. Polynomial is

$$p(\lambda) = (\lambda - \langle z, x \rangle)\lambda^{n-1}$$

In the case of orthogonal vectors w and z, then,

$$p(\lambda) = \lambda^n$$

If w and z are not orthogonal, even though there are only two fuzzy eigen values, it can be stated that the fuzzy eigenvalue 0 has a multiplicity of (n - 1).

Theorem – 3:

Each polynomial of degree n with a leading coefficient of 1, i.e.

$$q(\lambda) = \lambda^n + b_1\lambda^{n-1} + \dots + b_{n-1}\lambda + b_n$$

It serves as the characteristic polynomial for a certain fuzzy matrix. Proof:

Let us suppose a n × n fuzzy square matrix

$$B = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots \\ b_{n-1} & b_{n-2} & \dots & b_1 & b_n \end{pmatrix}$$

Then λI - B has the form

$$\lambda I - B = \begin{pmatrix} \lambda & -1 & 0 & \dots & 0 \\ 0 & \lambda & -1 & \dots & 0 \\ \dots & \dots & \lambda & \dots & \dots \\ b_{n-1} & b_{n-2} & \dots & \lambda + b_1 & b_n \end{pmatrix}$$

Next, perform a expansion in minor along the bottom row to obtain

$$\det(\lambda I - B) = b_n(-1)^{n+1} \det \begin{pmatrix} -1 & 0 & \dots & 0 \\ \lambda & -1 & \dots & 0 \\ \dots & \dots & \dots & \dots \end{pmatrix} + \dots + b_1(-1)^{n+n} \det \begin{pmatrix} \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots \\ \lambda & -1 & 0 & \dots \end{pmatrix}$$

$$= b_n(-1)^{n+1}(-1)^{n-1} + b_{n-1}(-1)^{n+2}\lambda(-1)^{n-2} + \dots + (\lambda + b_1)(-1)^{n+1}\lambda^{n-1} = b_n + b_{n-1}\lambda + \dots + b_1\lambda^n$$

Which is what we set out to prove.

Multiplicity of fuzzy matrix:

Let $A \in M_n(C)$ be a fuzzy matrix. Since $p(\lambda)$ represents a polynomial with a precise degree of n , therefore, it necessitates a precise count of n fuzzy eigenvectors $\lambda_1, \lambda_2, \dots, \lambda_n$.

The multiplicity of a fuzzy eigenvalue is determined by how often the monomial $(\lambda - \lambda_i)$ appears in the factorization of $p(\lambda)$.

Suppose $\mu_1, \mu_2, \dots, \mu_k$ are the distinct fuzzy eigen values with multiplicities m_1, m_2, \dots, m_k respectively. Then the ch. polynomial can be written as

$$p(\lambda) = \det(\lambda I - A) = \prod_{i=1}^n (\lambda - \lambda_i) = \prod_{i=1}^k (\lambda - \mu_i)^{m_i}$$

The multiplicities m_1, m_2, \dots, m_k are referred to as the fuzzy algebraic multiplicities associated with their corresponding fuzzy eigenvalues.

Algebraic multiplicity: The algebraic multiplicity of a fuzzy eigenvalue is the number of times it appears as a root of the characteristic polynomial (i.e., the polynomial whose roots are the fuzzy eigenvalues of a matrix).

Geometric multiplicity: The geometric multiplicity of a fuzzy eigenvalue is the dimension of the linear space of its associated fuzzy eigenvectors (i.e., its eigenspace).

Proposition: Let A be a $k \times k$ fuzzy matrix, $k \in [0,1]$. Let λ_k be one of the fuzzy eigenvalues of A . Then, the fuzzy geometric multiplicity of λ_k is less than or equal to its fuzzy algebraic multiplicity.

Proof: Suppose that the fuzzy geometric multiplicity of λ_k is equal to r , so that there is r linearly independent fuzzy eigenvectors x_1, x_2, \dots, x_r associated to λ_k . Arbitrarily choose $k-r$ vectors $x_{r+1}, x_{r+2}, \dots, x_k$, all having dimension $k \times 1$ and such that the n column vectors x_1, x_2, \dots, x_k are linearly independent. Define the $k \times k$ fuzzy square matrix

$$X = [x_1 \dots x_k]$$

For any r , denoted by b_r the vector that solves

$$X b_r = A x_r$$

which is guaranteed to exist because X is full-rank (its columns are linearly independent). Define the $k \times (k - r)$ fuzzy matrix

$$B = [b_{r+1} \dots b_k]$$

and denote by C its upper $r \times (k - r)$ block and by D its lower $(k - r) \times (k - r)$ block

$$B = \begin{pmatrix} C \\ D \end{pmatrix}$$

Denote by I_k the $k \times k$ fuzzy identity matrix. For any scalar λ , we have that

$$\begin{aligned} & (\lambda I_k - A)X \\ &= \begin{bmatrix} (\lambda I_k - A)x_1 & \dots & (\lambda I_k - A)x_r & (\lambda I_k - A)x_{r+1} & \dots & (\lambda I_k - A)x_k \end{bmatrix} \\ &= \begin{bmatrix} (\lambda - \lambda_k)x_1 & \dots & (\lambda - \lambda_k)x_r & \lambda x_{r+1} - A x_{r+1} & \dots & \lambda x_k - A x_k \end{bmatrix} \\ &= \begin{bmatrix} (\lambda - \lambda_k)x_1 & \dots & (\lambda - \lambda_k)x_r & \lambda x_{r+1} - X b_{r+1} & \dots & \lambda x_k - X b_k \end{bmatrix} \\ &= X \begin{bmatrix} (\lambda - \lambda_k)I_r & -C \\ 0 & \lambda I_{k-r} - D \end{bmatrix} \end{aligned}$$

Since X is full-rank and, as a consequence its determinant is non-zero, we can write

$$\begin{aligned}
 & |\lambda I_K - A| \\
 &= |\lambda I_K - A| \cdot |X| \cdot \frac{1}{|X|} \\
 &= |(\lambda I_K - A)X| \cdot \frac{1}{|X|} \\
 &= \left| X \begin{bmatrix} (\lambda - \lambda_k)I_r & -C \\ 0 & \lambda I_{K-r} - D \end{bmatrix} \right| \cdot \frac{1}{|X|} \\
 &= |X| \left| \begin{bmatrix} (\lambda - \lambda_k)I_r & -C \\ 0 & \lambda I_{K-r} - D \end{bmatrix} \right| \cdot \frac{1}{|X|} \\
 \boxed{A} &= |(\lambda - \lambda_k)I_r| \cdot |\lambda I_{K-r} - D| \\
 &= (\lambda - \lambda_k)^r |\lambda I_{K-r} - D|
 \end{aligned}$$

where in the last step we have used a result about the determinant of block-matrices. The fuzzy eigenvalues of A solve the characteristic equation

$$|\lambda I_K - A| = 0$$

or, equivalently, the equation

This equation has a root

$$(\lambda - \lambda_k)^r |\lambda I_{K-r} - D| = 0$$

$$\lambda = \lambda_k$$

that is repeated at least r times. Therefore, the fuzzy algebraic multiplicity of λ_k is at least equal to its fuzzy geometric multiplicity r. It can be larger if λ_k is also a root of

$$|\lambda I_{K-r} - D| = 0$$

3. RESULTS

This paper provides an insight into the concept of fuzzy metric spaces with a focus on fuzzy eigenvalues and fuzzy eigenvectors. We present several theorems concerning the eigenvalues and eigenvectors within the context of a fuzzy metric space. Notably, we find that many propositions pertaining to metric spaces are applicable to fuzzy eigenvalues and eigenvectors. Additionally, we attempt to offer some insights into the multiplicity of fuzzy matrices. This field presents significant opportunities for further exploration by researchers, indicating the vast potential for future study and development.

CONFLICT OF INTEREST

The authors state that they do not have any conflicts of interest.

REFERENCES

[1] I. Kramosil, J. Michálek, "Fuzzy metric and statistical metric spaces," *Kybemetika*, 11(1975), pp.336-344
 [2] M. Grabiec, "Fixed points in fuzzy metric spaces," *Fuzzy sets and systems*, 27(1988), pp.385-389
 [3] L.Boxer, "On Hausdorff-like metric for fuzzy sets," *Pattern Recognition Letters* 18 (1997), 115-118.
 [4] B. B. Chaudhuri and A. Rosenfeld, "On a metric distance between fuzzy sets," *Pattern Recognition*

Letters 17 (1996), 1157-1160.
 [5] P. Diamond and P. Kloden, "Metric spaces of fuzzy sets," *Fuzzy Sets and Systems* 35 (1990), 241-249
 [6] O. Kaleva and S. Seikkala, "On fuzzy metric space," *Fuzzy Sets and Systems* 12(1984), 215-229.
 [7] C. Felbin, "Finite dimensional fuzzy normed linear space," *Fuzzy Sets and Systems* 48 (1992), 239-248.
 [8] A. George and P. V. Veeramani, "On some results of fuzzy metric spaces," *Fuzzy Sets and Systems* 64 (1994), 395-399.
 [9] A. George and P. V. Veeramani, "On some results of analysis for fuzzy metric spaces," *Fuzzy Sets and Systems* 90 (1997), 365-368.
 [10] V. Gregori and S. Romaguera, "Some properties of fuzzy metric spaces," *Fuzzy Sets and Systems* 115 (2000), 485-489.
 [11] O. Hadzic and E. Pap, "A fixed point theorem for multivalued mapping in probabilistic metric spaces and an application in fuzzy metric spaces," *Fuzzy Sets and Systems* 127 (2002), 333-344.
 [12] J. L. Fan, "Note on Hausdorff-like metrics for fuzzy sets," *Pattern Recognition Letters* 19 (1998), 793-796.
 [13] P. Brass, "On the no existence of Hausdorff-like metrics for fuzzy sets," *Pattern Recognition Letters* 23 (2002), 39-43.
 [14] G. Przemyslaw, "Metrics and orders in space of fuzzy

- numbers,” *Fuzzy Sets and Systems* 97 (1998), 83-94.
- [15] W. Voxman, “Some remarks on distance between fuzzy numbers,” *Fuzzy Sets and Systems* 100 (1998), 353-365.
- [16] Zun-Quan Xia and Fang-Fang Guo, “Fuzzy Metric Spaces,” *J. Appl. Math. & Computing* Vol. 16(2004), No. 1 - 2, pp. 371 – 381.
- [17] P. Priya, R. Ramya, S. Sangeetha, A. Saranya, “Basic Concepts of Fuzzy Matrices”, *International Journal of Engineering Research and Technology*, Volume 08, Issue 12 (December 2019).
- [18] W. Kolodziejczyk, convergence of powers of s.transitive fuzzy matrices, *fuzzy sets and systems*; 26(1988),127-130.
- [19] L.J. Xin, controllable fuzzy matrices, *fuzzy sets and systems*,45(1992), 313-319.
- [20] L.J. Xin, convergence of powers of controllable fuzzy matrices, *fuzzy sets and systems*, 62(1994), 83-88.
- [21] M.Z. Ragab and E.G. Emam, The determinant and adjoint of a square fuzzy matrix, *fuzzy sets and systems*, 61(1994),297-307.
- [22] J.B. Kim, Determinant theory for fuzzy and Boolean matrices, *Congressus Numerantium*, (1988),273-276.
- [23] A.K. Shyamal and M. Pal, Two new operators on fuzzy matrices, *J.Applied mathematics and computing*, 15(2004), 91-107.
- [24] A.K. Shyamal and M. Pal, Distance between fuzzy matrices its applications, *ActasienciaIndica*, XXXF.M(1) (2005),199-204
- [25] Michael G.R. Voskoglou, Applications of Fuzzy linear programming to Business problems, *WSEAS Transactions on Mathematics*,2020,Vol.19,pp.343 – 348
2.[URL: <https://www.wseas.org/multimedia/journals/mathematics/2020/a685106-050.pdf>]
- [26] R. Ghanbari et al., Fuzzy linear programming problems: models and solutions, *Soft Computing*, 2020, 24: pp.10043 –10073 [DOI: <https://doi.org/10.1007/s00500-019-04519-w>]
- [27] Ebrahimnejad and Verdegay, Fuzzy sets-Based methods and techniques for modern analytics, *Studies in Fuzziness and Soft computing*. Springer, 2018, Vol 364.
- [28] [DOI: <https://doi.org/10.1007/978-3-319-73903-8>]
- [29] Samsun Nahar et al., Solving Fuzzy linear programming problem using weighted sum and comparisons with ranking function, *International Journal of Scientific & Engineering Research*, 2019, Vol.10, Issue 11, pp.480 -484.