

The Interplay of Chromatic Contrast and Typographic Legibility: Enhancing Mobile Interface Accessibility for Seniors with Age-Related Visual and Cognitive Declines

Bhaskar Mani Tiwari,

P.hD Scholar, Deptt. of Computer Science and IT, IIS University, Jaipur

Prof. Swati V Chande,

H.O.D, Deptt. of Computer Science and IT, IIIM, Jaipur

Abstract

Age-related visual and cognitive impairments significantly affect seniors' ability to interact with mobile interfaces, yet many contemporary applications fail to address these accessibility challenges adequately. This research examines the synergistic relationship between chromatic contrast and typographic legibility in mobile interface design for older adults aged 65 and above. Through a comprehensive analysis of current accessibility standards, empirical studies, and age-related physiological changes, this paper establishes evidence-based design guidelines that accommodate declining contrast sensitivity, reduced visual acuity, and cognitive processing limitations. The findings reveal that implementing WCAG 2.2 Level AAA contrast ratios (7:1 for normal text) combined with font sizes of 18 points or larger significantly enhances readability and reduces visual fatigue for seniors. Furthermore, the research demonstrates that carefully selected color combinations particularly high-contrast schemes such as black text on white backgrounds combined with sans-serif typefaces can compensate for age-related visual processing deficits. This paper provides practical design recommendations for creating inclusive mobile interfaces that bridge the digital divide and promote technological independence among the aging population.

Keywords: Mobile accessibility, chromatic contrast, typographic legibility, age-related visual decline, senior interface design

1. Introduction

The global demographic shift towards an aging population presents unprecedented challenges for mobile interface design, with adults aged 65 and older representing the fastest-growing user demographic worldwide. As mobile technologies increasingly mediate essential services including healthcare management, financial transactions, and social connectivity the accessibility of these interfaces becomes critical to maintaining seniors' independence and quality of life. However, age-related physiological changes create substantial barriers to mobile technology adoption, with over 50% of individuals aged 65 and above experiencing some form of visual impairment that directly affects their ability to interact with digital interfaces.

Age-related visual declines manifest through multiple mechanisms: reduced contrast sensitivity, decreased visual acuity, diminished color discrimination, and increased susceptibility to glare. Simultaneously, cognitive changes including slower processing speeds, reduced working memory capacity, and decreased attention span compound these visual challenges, creating a multifaceted accessibility problem that cannot be addressed through single-dimension solutions. Despite these well-documented challenges, many mobile applications prioritize aesthetic minimalism over functional accessibility, employing low-contrast color schemes and small typography that render interfaces effectively unusable for older adults.

The interplay between chromatic contrast and typographic legibility represents a critical yet underexplored dimension of age-friendly mobile design.

While existing research has examined these factors independently, their synergistic effects on comprehension, task completion, and user satisfaction among seniors remain inadequately understood. This research gap is particularly problematic given that color and typography do not function as isolated variables but interact dynamically to influence legibility, readability, and cognitive load.

This paper addresses this gap by systematically examining how chromatic contrast and typographic variables collectively impact mobile interface accessibility for seniors with age-related visual and cognitive declines. The research synthesizes findings from gerontology, human-computer interaction, visual psychophysics, and accessibility standards to develop evidence-based design guidelines that accommodate the specific needs of aging users.

2. Literature Review

2.1 Age-Related Visual Decline and Mobile Interface Interaction

Aging fundamentally alters the visual system through multiple physiological pathways that collectively degrade perceptual capacity. Dobres et al. (2016) demonstrated that older adults experience significant reductions in sensitivity to high spatial frequencies, resulting in perceived blur that disproportionately affects their ability to recognize fine textual details. Their psychophysical experiments revealed that performance thresholds for text legibility shifted significantly between younger and older cohorts, with older observers' performance declining at substantially lower blur levels (1.04 versus 2.16 arcmin at the 90% threshold).

Contrast sensitivity the ability to distinguish objects from their backgrounds diminishes progressively with age, creating particular challenges for mobile interfaces that often employ subtle color gradations for aesthetic purposes. Research indicates that visual acuity declines correlate strongly with age, even exceeding the correlation between cloudiness or specific pathologies and reading performance. This age-dependent degradation affects not only text recognition but also button identification, icon comprehension, and overall navigation efficiency.

Kobayashi et al. (2011) conducted comprehensive usability evaluations with elderly participants using mobile touchscreen interfaces, identifying that while touchscreens were generally accessible, careful

observation revealed systematic problems stemming from hard-to-see displays and difficulty distinguishing interactive elements. Their longitudinal study demonstrated that while experience improved proficiency over one week, fundamental visual limitations persisted, suggesting that interface modifications rather than user training represent the more effective intervention strategy.

2.2 Color Perception and Chromatic Contrast in Aging Populations

Age-related changes in color perception significantly impact mobile interface usability, particularly for users with age-related macular degeneration (AMD) and other degenerative conditions. Recent studies using smartphone-based color testing have revealed that AMD patients exhibit significant impairments in color contrast discrimination, particularly for blue and yellow wavelengths, while red-green discrimination remains relatively preserved. These findings have important implications for mobile color scheme selection, as they indicate that commonly used blue interface elements may be particularly problematic for older users with AMD.

The Web Content Accessibility Guidelines (WCAG) 2.2 establish specific contrast ratio requirements based on age-related visual capabilities. The Level AA standard of 4.5:1 for normal text was specifically calibrated to compensate for contrast sensitivity loss equivalent to approximately 20/40 vision, which represents typical visual acuity for individuals around age 80. The more stringent Level AAA requirement of 7:1 accommodates users with approximately 20/80 vision, providing additional compensation for both low vision and color deficiency.

Research on high-contrast color schemes consistently demonstrates that older users benefit from stark differentiation between foreground and background elements. Studies indicate that 70% of older users experience vision-related difficulties that are ameliorated by contrast ratios of at least 4.5:1, with many researchers recommending higher ratios for critical interface elements. Dark text on light backgrounds or vice versa represents the most effective approach, as these combinations minimize the visual processing demands imposed on aging visual systems.

2.3 Typographic Legibility and Font Selection for Seniors

Font size represents one of the most critical variables affecting mobile interface accessibility for older adults, yet optimal specifications remain contested in the literature. Recent systematic reviews examining reading time, readability, and legibility for older adults consistently recommend font sizes of 18 points as significantly faster to read than smaller alternatives. Experimental comparisons of 14, 16, and 18-point fonts across both serif and sans-serif typefaces revealed that 18 points reduced visual fatigue during both detailed and skim reading tasks.

However, font size alone provides an incomplete solution, as typeface characteristics substantially modulate legibility effects. Research comparing different typefaces found that older participants demonstrated greater sensitivity to font legibility differences than younger readers, with performance varying significantly based on reading scenario demands. For critical print size tasks requiring recognition at small sizes, older participants achieved better performance with certain typefaces (KBH Text) compared to aesthetically preferred alternatives (Gill Sans).

The serif versus sans-serif debate remains unresolved, with cultural and contextual factors mediating preferences. UK participants in cross-cultural studies preferred 18-point sans-serif fonts, finding them easier and less tiring to read, while Thai participants showed opposite preferences for serif fonts. These findings suggest that optimal typography may require cultural adaptation rather than universal specifications.

Character spacing, stroke weight, and letter width interact with age-related visual processing to affect reading speed and comprehension. As individuals age, they demonstrate reduced ability to shorten viewing distance to compensate for small character sizes a strategy younger users employ effectively likely due to presbyopia and reduced accommodative flexibility.

2.4 Cognitive Load and Interface Complexity

Age-related cognitive decline compounds visual impairments, creating multilayered accessibility challenges that demand integrated design solutions. Czaja and Sharit's extensive research through the CREATE Center demonstrates that older adults exhibit willingness to adopt mobile technologies when interfaces minimize cognitive load through clear navigational

structures, consistent interaction patterns, and reduced information density.

Design guidelines for mobile health applications targeting seniors emphasize the importance of consistent interfaces that match the cognitive processing capabilities of older adults. Three key principles emerge across multiple studies: simplified navigation, reduced cognitive load through clear hierarchical structure, and interface consistency that supports the development of mental models. The interaction between cognitive processing demands and visual perception creates compounding effects, where users experiencing cognitive strain demonstrate even greater difficulty distinguishing interactive from non-interactive elements.

2.5 Comprehensive Design Guidelines for Senior-Friendly Mobile Interfaces

Amouzadeh et al. (2025) synthesized design principles from multiple disciplines, emphasizing that age-friendly interfaces must simultaneously address sensory limitations, reduced contrast sensitivity, motor skill decline, and cognitive processing changes. Their systematic analysis revealed that many mobile interfaces fail to consider these common age-related problems, resulting in unnecessarily high abandonment rates and reduced technology adoption among seniors.

A landmark systematic review by Silva et al. (2023) analyzed 40 empirical studies conducting usability tests with populations aged over 60 years, extracting 27 evidence-based design guidelines organized into five categories: Help & Training, Navigation, Visual Design, Cognitive Load, and Interaction. Two principles emerged as transversal and of greatest significance: "Simplify" and "Increase the size and distance between interactive controls". Within the Visual Design category, specific recommendations addressed layout, icons, and appearance, with particular emphasis on providing high contrast between foreground and background colors.

3. Conceptual Framework and Design Principles

3.1 Age-Related Visual and Cognitive Changes

Mobile interface accessibility for seniors must be understood through the lens of multiple, concurrent physiological changes affecting both visual perception and cognitive processing. These changes are not discrete impairments but rather interconnected alterations that synergistically impact the user experience.

Visual System Changes:

- Reduced contrast sensitivity affecting object-background discrimination
- Decreased visual acuity and sensitivity to high spatial frequencies
- Lens yellowing and increased light scatter causing glare sensitivity
- Impaired color discrimination, particularly for blue-yellow spectra
- Reduced accommodative flexibility (presbyopia) limiting focus adjustment
- Slower dark adaptation and reduced peripheral vision

Cognitive Processing Changes:

- Reduced working memory capacity affecting multitasking and information retention
- Slower processing speeds impacting reaction time and decision-making
- Decreased attention span and increased susceptibility to distraction
- Greater difficulty forming mental models of novel interfaces
- Reduced ability to recover from errors and navigate complex hierarchies

These changes necessitate design approaches that reduce both perceptual demands and cognitive load simultaneously, as compensatory strategies that address one dimension may inadvertently increase demands in another.

3.2 The Synergistic Relationship Between Contrast and Typography

Chromatic contrast and typographic legibility do not function independently but interact to determine overall interface accessibility. High contrast alone cannot compensate for inadequate font sizing, nor can large typography overcome insufficient color differentiation. The optimal design space emerges at the intersection of these variables, where both dimensions meet or exceed age-appropriate thresholds.




Figure 1. Conceptual interaction model of chromatic contrast and typographic size on legibility for different age groups

Source: (adapted and synthesized from): Silva et al. (2023) and Huang et al. (2022).

This figure illustrates the joint influence of chromatic contrast ratio and font size on perceived legibility. Two conceptual surfaces represent younger users and older users (65+). The surface corresponding to older adults is shifted towards higher contrast ratios and larger font sizes, indicating that seniors require simultaneously higher contrast and larger typography to reach comparable levels of legibility.

The model highlights the synergistic relationship between contrast and typographic size: improving one parameter alone is insufficient to achieve optimal legibility for older users when the other parameter remains below age-appropriate thresholds.

3.3 WCAG Standards and Senior-Specific Requirements

The WCAG 2.2 guidelines provide foundational accessibility requirements, but their minimum standards may prove insufficient for users with significant age-related impairments. While Level AA compliance (4.5:1 contrast ratio) addresses typical age 80 visual acuity, optimal senior accessibility often requires Level AAA standards (7:1 contrast ratio).

Table 1: WCAG Levels

WCA G Level	Contra st Ratio	Visual Acuity Equivale nt	Recommend ed Application
Level AA	4.5:1 (normal text)	20/40 vision (age ~80)	Minimum standard for general senior population
Level AA	3:1 (large text ≥18pt)	Same as above	Acceptable for headings and large UI elements
Level AAA	7:1 (normal text)	20/80 vision	Recommend ed for critical information and users with significant impairment
Level AAA	4.5:1 (large text ≥18pt)	Same as above	Recommend ed for all large text elements

4. Design Recommendations and Best Practices

4.1 Chromatic Contrast Specifications

Primary Text Elements:

Research consistently demonstrates that seniors benefit most from maximum contrast configurations. Black text (#000000) on white backgrounds (#FFFFFF) provides a 21:1 contrast ratio, substantially exceeding WCAG AAA requirements and accommodating users with severe visual impairments. Alternatively, dark gray (#333333) on off-white (#F8F8F8) offers a slightly softer 13.5:1 ratio that reduces glare while maintaining excellent legibility.

Secondary and Tertiary Elements:

Interactive elements, labels, and secondary information should maintain minimum 7:1 contrast ratios. User interface components such as buttons, form fields, and navigation elements must meet the WCAG 2.2 requirement of 3:1 contrast against adjacent colors.

Color Combinations to Avoid:

Given research on AMD-related color perception deficits, designers should exercise caution with blue interface elements, particularly against purple or green backgrounds. Low-contrast pairings such as gray text on gray backgrounds, yellow text on white, or light blue on white systematically fail to meet senior accessibility needs.

4.2 Typographic Specifications

Font Size:

Body text should employ 18-point fonts as the baseline, with 16 points as the absolute minimum for secondary information. Headings and critical labels should scale proportionally, typically 24-32 points depending on hierarchical level. Research demonstrates that 18-point fonts significantly reduce reading time and visual fatigue compared to smaller alternatives.

Typeface Selection:

Sans-serif typefaces generally provide superior legibility for screen-based reading among Western seniors, though cultural factors may influence preferences. Recommended typefaces include Arial, Helvetica, Verdana, and Roboto, all designed with clear character differentiation and moderate x-heights. Avoid decorative, script, or highly condensed typefaces that increase character recognition demands.

Character and Line Spacing:

Line height should be set to 1.5-1.8 times the font size to prevent text crowding and support tracking across lines. Letter spacing (tracking) should be slightly increased approximately 0.05-0.1em to enhance character differentiation for users with reduced spatial frequency sensitivity.




Figure 2. Comparison of optimal and sub-optimal typographic configurations for senior users

Source (based on): Amouzadeh et al. (2025) and Silva et al. (2023).

This figure presents two representative text samples illustrating the effect of typographic configuration and chromatic contrast on readability for older adults.

Panel A (Optimised configuration) demonstrates a senior-friendly text layout using an 18-point sans-serif typeface, high foreground-background contrast (approximately 7:1), and increased line spacing (approximately 1.5). This configuration supports faster visual parsing, improved character discrimination, and reduced visual fatigue for older users.

Panel B (Sub-optimal configuration) shows a commonly observed mobile interface layout using a 12-point serif typeface, low contrast (approximately 3:1), and tight line spacing (approximately 1.0). This combination increases visual crowding and reduces legibility, particularly for users with age-related declines in contrast sensitivity and spatial frequency processing.

Readability metrics displayed in the figure

- Panel A: 18 pt, sans-serif, $\approx 7:1$ contrast, line spacing 1.5
- Panel B: 12 pt, serif, $\approx 3:1$ contrast, line spacing 1.0

4.3 Interactive Element Design

Button and Touch Target Sizing:

Touch targets should measure minimum 44×44 pixels (approximately 9-10mm), exceeding standard recommendations to accommodate reduced motor precision. Research demonstrates that increasing touch target sizes significantly reduces error rates and user frustration among elderly users.

Visual Affordances:

Interactive elements must be clearly distinguishable from static content through explicit visual cues including borders, shadows, and color differentiation. Users experiencing cognitive strain demonstrate particular difficulty identifying interactive elements when conventional affordances are absent.

4.4 Layout and Information Architecture

Simplification and Focus:

Interfaces should present single primary tasks per screen, minimizing cognitive load and reducing working memory demands. Information density should be substantially reduced compared to interfaces targeting

younger demographics, with generous whitespace supporting visual parsing.

Consistency and Predictability:

Navigation patterns, interaction paradigms, and visual hierarchies must remain consistent throughout the application to support mental model formation. Older adults benefit particularly from predictable layouts that reduce exploratory demands and support recognition over recall.

Table 2: Comparison of Standard vs. Senior-Optimized Mobile Interface Specifications

Design Element	Standard Mobile Practice	Senior-Optimized Specification	Evidence Base
Body text size	12-14pt	18pt minimum	Huang et al. (2022)
Text contrast ratio	4.5:1 (AA)	7:1 (AAA)	WCAG 2.2 (2024)
Touch target size	44×44px	48×48px minimum with spacing	Silva et al. (2023)
Line height	1.2-1.4×	1.5-1.8×	Silva et al. (2023)
Information density	Medium-high	Low with generous whitespace	Amouzadeh et al. (2025)
Color palette	Varied, often low-contrast	High-contrast, limited palette	Multiple sources

5. Methodology for Evaluating Interface Accessibility

5.1 Assessment Framework

Comprehensive evaluation of mobile interface accessibility for seniors requires multi-method

assessment combining objective measurements with subjective user feedback. The evaluation framework should incorporate:

Automated Contrast Analysis:

Tools such as AccessibilityScanner (Android) and Contrast Checker (WebAIM) provide quantitative contrast ratio measurements against WCAG standards. However, automated tools cannot capture contextual factors affecting perceived legibility, necessitating supplementary methods.

Legibility Testing with Age-Representative Samples:

Empirical testing must include participants aged 65+ representing the target demographic, ideally stratified by visual acuity levels and cognitive function. Testing protocols should measure reading speed, comprehension accuracy, task completion rates, and subjective fatigue across multiple sessions to account for learning effects.

Psychophysical Measurements:

Following methodologies established by Dobres et al. (2016), psychophysical techniques can quantify legibility thresholds, determining the minimum contrast ratios and font sizes at which older adults maintain 90% accuracy. These measurements provide objective baselines for design specifications.

5.2 User Testing Protocols

Usability testing with seniors should employ age-appropriate methodologies recognizing potential fatigue, reduced attention span, and unfamiliarity with testing environments. Sessions should be limited to 45-60 minutes with regular breaks, use simplified think-aloud protocols, and provide extensive practice opportunities before data collection.

Tasks should reflect authentic use cases relevant to older adults' lives, such as medication management, appointment scheduling, or family communication, rather than abstract or gamified scenarios that may lack motivational salience. Kobayashi et al. (2011) demonstrated that longitudinal evaluations testing users after one week of experience reveal persistent problems that initial testing might attribute to unfamiliarity.




Figure 3. Comprehensive evaluation methodology for senior-friendly mobile interfaces

Source (methodological framework synthesised from): Dobres et al. (2016), Kobayashi et al. (2011) and Silva et al. (2023).

This figure illustrates a multi-track evaluation framework for assessing mobile interface accessibility for older adults.

Three parallel assessment streams are shown:

- Automated accessibility testing, including contrast and interface compliance checking using accessibility analysis tools.
- Psychophysical measurements, focusing on objective legibility thresholds and accuracy levels under controlled visual conditions.
- User testing with older adults (65+), measuring reading speed, task completion performance and subjective visual fatigue.

All three evaluation streams converge into an integrated accessibility assessment and design refinement stage, where objective measurements and user-centred findings are jointly used to inform iterative interface improvements.

The framework highlights that reliable accessibility evaluation for seniors requires the combination of automated inspection, experimental visual performance testing and real-world usability assessment.

6. Discussion

6.1 Integrating Contrast and Typography: Beyond Additive Effects

The relationship between chromatic contrast and typographic legibility transcends simple additive models, exhibiting interaction effects that demand holistic design approaches. Research demonstrates that inadequate contrast disproportionately impacts small typography, while large fonts provide limited benefit when presented with insufficient color differentiation. This synergy

suggests that designers must optimize both dimensions simultaneously rather than treating them as independent variables subject to individual threshold requirements.

The WCAG guidelines acknowledge this interaction through differentiated standards for normal versus large text, reducing contrast requirements from 4.5:1 to 3:1 when font size reaches 18 points. However, for seniors with significant visual impairments, even large text benefits from maximum contrast, as age-related changes in spatial frequency sensitivity affect text perception across all sizes.

6.2 Cultural and Contextual Considerations

The cross-cultural research on font preferences reveals that optimal typography cannot be universally prescribed but must accommodate cultural reading habits and linguistic characteristics. The divergent preferences between UK participants favoring sans-serif and Thai participants preferring serif fonts suggest that character systems, reading direction, and cultural aesthetic norms mediate legibility effects. This finding has important implications for global mobile applications serving diverse senior populations.

Environmental context also substantially affects contrast requirements, with outdoor usage, varying ambient lighting, and screen brightness limitations creating dynamic accessibility challenges. Adaptive interfaces that adjust contrast ratios and font sizes based on environmental sensors represent a promising direction for enhancing real-world usability beyond laboratory-optimized configurations.

6.3 Balancing Accessibility and Aesthetics

Design practitioners often express concerns that high-contrast, large-typography interfaces appear dated or aesthetically compromised compared to minimalist contemporary designs. However, this perspective reflects ageist assumptions that older adults prioritize aesthetics over functionality. Research by Czaja and Sharit demonstrates that seniors evaluate technologies primarily based on perceived usefulness and ease of use rather than visual sophistication. Interfaces that enable successful task completion and reduce frustration cultivate greater user satisfaction than aesthetically refined but functionally inaccessible alternatives.

Moreover, accessibility features benefit broader populations beyond their intended demographic a phenomenon known as the "curb-cut effect". High-contrast interfaces assist users in bright sunlight, large

typography benefits users with temporary visual impairments or fatigue, and simplified navigation aids users under cognitive load regardless of age. Designing for seniors thus represents not a niche accommodation but a universal design approach enhancing usability across diverse contexts and user characteristics.

6.4 Technology Adoption and Digital Inclusion

The systematic accessibility barriers in mobile interfaces contribute substantially to the digital divide separating seniors from essential services and social participation. As healthcare systems, financial services, and government agencies increasingly mandate digital interactions, inaccessible interfaces effectively exclude older adults from full civic and economic participation. This exclusion carries serious consequences for health outcomes, financial security, and social connectivity, particularly for socioeconomically disadvantaged seniors lacking alternative access channels.

Research from the CREATE Center demonstrates that technophobic stereotypes do not reflect empirical reality older adults exhibit strong willingness to adopt technologies when interfaces accommodate their capabilities and the value proposition is clear. Czaja reports successfully training adults up to 98 years old in computer use, demonstrating that motivation and appropriate design overcome age-related limitations. Interface accessibility thus represents not a concession to inevitable decline but an enabling factor unlocking older adults' technological engagement.

7. Implications for Design Practice

7.1 Immediate Implementation Priorities

Practitioners developing or redesigning mobile applications for older adults should prioritize:

1. **Audit existing interfaces** using automated contrast checking tools and identify all elements failing WCAG AA standards
2. **Implement minimum 18-point font sizes** for all body text, with proportional increases for headings and critical labels
3. **Adopt high-contrast color schemes** using black or dark gray text on white or light backgrounds, achieving minimum 7:1 ratios
4. **Increase touch target sizes** to 48×48 pixels minimum, with adequate spacing preventing accidental activation

5. **Simplify navigation** to single-level hierarchies where possible, with clear labeling and consistent interaction patterns

7.2 User-Centered Design and Co-Design Approaches

Following recommendations by Mazuz and Biswas, participatory design approaches involving older adults throughout the development lifecycle yield substantially more usable outcomes than designer-driven specification. Co-design sessions enable designers to observe authentic interaction patterns, identify unanticipated barriers, and validate design decisions against real-world usage rather than assumptions about senior capabilities.

Crucially, co-design participants should represent the diversity of the senior population, including individuals with varying visual acuities, cognitive function levels, technology experience, and socioeconomic backgrounds. Recruitment limited to tech-savvy "super seniors" produces unrepresentative feedback that may not generalize to broader populations experiencing significant age-related declines.

7.3 Adaptive and Customizable Interfaces

Given the heterogeneity of aging experiences, interfaces offering customization options enable users to configure optimal settings matching their individual capabilities. Preferences for contrast themes (including dark mode), font size adjustment, and spacing modifications should be easily accessible and persist across sessions. However, customization options must be presented clearly with preview functionality, as complex configuration interfaces may themselves pose accessibility barriers.

8. Limitations and Future Research Directions

8.1 Research Limitations

Current research on senior mobile accessibility exhibits several limitations constraining generalizability. Most studies employ laboratory settings with controlled lighting and dedicated attention, potentially overestimating real-world usability where environmental distractions, variable lighting, and multitasking demands affect performance. Longitudinal studies tracking technology adoption and sustained usage patterns remain scarce, limiting understanding of how initial usability translates to long-term engagement.

Sample sizes in usability studies frequently remain modest, and recruitment often skews toward relatively

healthy, cognitively intact seniors, underrepresenting individuals with significant impairments who face the greatest accessibility barriers. Cross-cultural research remains limited, with Western populations dominating empirical studies despite substantial global variation in aging demographics, technology access, and design preferences.

8.2 Future Research Priorities

Several critical research directions warrant investigation:

Longitudinal Adoption Studies: Research examining sustained mobile technology usage among seniors over extended periods (6-12 months) would illuminate how initial usability affects long-term adoption, identify evolving accessibility needs as users age, and reveal whether initial interface barriers or ongoing usability problems drive abandonment.

Cognitive Decline Trajectories: Studies differentiating accessibility requirements across cognitive decline stages from healthy aging through mild cognitive impairment to dementia would enable more nuanced design guidelines tailored to specific populations. Research with Alzheimer's patients demonstrates substantial performance benefits from adapted interfaces, but optimal specifications remain inadequately characterized.

Multimodal Interaction Design: Investigation of voice-activated features, haptic feedback, and multimodal interfaces combining visual, auditory, and tactile channels may compensate for visual limitations through alternative sensory pathways. Research should examine whether multimodal approaches reduce cognitive load or inadvertently increase complexity through unfamiliar interaction paradigms.

Artificial Intelligence and Adaptive Systems: Machine learning approaches enabling real-time interface adaptation based on user performance patterns, environmental conditions, and physiological measurements (e.g., eye tracking, interaction latency) represent promising frontiers. However, empirical validation with older adult populations remains limited.

Socioeconomic and Cultural Diversity: Research explicitly examining how socioeconomic status, educational background, cultural context, and prior technology experience moderate optimal design specifications would support more inclusive and equitable interface development.

9. Conclusion

The interplay of chromatic contrast and typographic legibility represents a critical determinant of mobile interface accessibility for seniors experiencing age-related visual and cognitive declines. This research demonstrates that optimal accessibility emerges not from isolated optimizations of individual design variables but from holistic approaches recognizing the synergistic relationships between color, typography, layout, and interaction design. Evidence consistently supports implementing WCAG Level AAA contrast standards (7:1 for normal text), employing minimum 18-point sans-serif typography, maximizing touch target sizes, and simplifying information architecture to accommodate the multifaceted challenges of aging visual and cognitive systems.

Current mobile interface practices systematically fail to meet the accessibility needs of the rapidly growing senior demographic, contributing to digital exclusion with serious consequences for health, economic security, and social participation. Yet research from Czaja, Sharit, and colleagues demonstrates that accessibility barriers reflect design choices rather than insurmountable limitations of aging users seniors exhibit strong willingness and capacity to adopt technologies when interfaces accommodate their capabilities. The persistent myth of the technophobic older adult thus obscures the reality that technology rejection reflects rational responses to poorly designed, inaccessible interfaces.

As populations age globally and digital technologies increasingly mediate essential services, inclusive design transcends ethical obligation to become functional necessity. The design principles articulated in this research grounded in empirical evidence from psychophysics, gerontology, and human-computer interaction provide actionable guidelines enabling practitioners to create mobile interfaces that genuinely serve users across the lifespan. Future research must extend these foundations through longitudinal adoption studies, investigations across cognitive decline trajectories, and examination of adaptive multimodal interfaces, while prioritizing participant diversity to ensure findings generalize across the heterogeneous senior population.

Ultimately, accessibility for seniors represents not specialized accommodation but fundamental usability. The "curb-cut effect" demonstrates that designing for users with the most significant needs produces interfaces

beneficial across diverse populations and contexts. By embracing the design challenge of age-related decline, the field advances toward truly universal mobile interfaces accessible, usable, and dignifying for all users regardless of age, ability, or circumstance.

References

1. Amouzadeh, E., et al. (2025) 'Optimizing mobile app design for older adults', *PMC*, 13 August. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC12350549/> (Accessed: 5 February 2026).
2. Czaja, S.J. and Sharit, J. (2017) 'Adapting for age: Research on aging and technology use', *APA Monitor on Psychology*, September. Available at: <https://www.apa.org/monitor/2017/09/adapting-age> (Accessed: 5 February 2026).
3. Dobres, J., et al. (2016) 'Age-related differences in the legibility of degraded text', *PMC*, 11 December. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC5256463/> (Accessed: 5 February 2026).
4. Dobres, J., et al. (2016) 'Utilising psychophysical techniques to investigate the effects of age, typeface design, and text size on legibility', *PMC*, 3 March. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC5213401/> (Accessed: 5 February 2026).
5. Huang, L., et al. (2022) 'How to design font size for older adults', *PMC*, 31 July. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC9376262/> (Accessed: 5 February 2026).
6. Kobayashi, M., et al. (2011) 'Elderly user evaluation of mobile touchscreen interactions', in *INTERACT 2011 Conference Proceedings*. Available at: <https://opendl.ifip-tc6.org/db/conf/interact/interact2011-1/> (Accessed: 5 February 2026).
7. Leat, S.J., et al. (2020) 'Use of the King-Devick Variable Color Contrast Sensitivity Chart to differentiate stages of age-related macular degeneration', *PMC*, 11 May. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC7228553/> (Accessed: 5 February 2026).
8. Munsamy, A.J., et al. (2022) 'A novel smartphone-based color test for detection of color vision defects in age related macular degeneration', *Wiley Online Library*. Available at: <https://onlinelibrary.wiley.com/doi/10.1155/2022/9744065> (Accessed: 5 February 2026).

9. Petersen, E.J. and Sherry, L. (2023) 'The effect of age and font on reading ability', *University of Cincinnati Journals*, 5(2). Available at: <https://journals.uc.edu/index.php/vl/article/view/4654> (Accessed: 5 February 2026).
10. Silva, P.A., et al. (2023) 'Design guidelines of mobile apps for older adults: Systematic review', *JMIR mHealth and uHealth*, 20 September. Available at: <https://mhealth.jmir.org/2023/1/e43186> (Accessed: 5 February 2026).
11. Vaportzis, E., et al. (2017) 'Design considerations for mobile health applications targeting older adults', *PMC*, 2 April 2014. Available at: <https://pmc.ncbi.nlm.nih.gov/articles/PMC8837196/> (Accessed: 5 February 2026).
12. W3C (2024) *Web Content Accessibility Guidelines (WCAG) 2.2*, 11 December. Available at: <https://www.w3.org/TR/WCAG22/> (Accessed: 5 February 2026).
13. W3C (2011) 'Understanding Success Criterion 1.4.3: Contrast (Minimum)', *Web Accessibility Initiative*, 31 March. Available at: <https://www.w3.org/WAI/WCAG22/Understanding/contrast-minimum.html> (Accessed: 5 February 2026).
14. WebAIM (2024) *Contrast Checker*. Available at: <https://webaim.org/resources/contrastchecker/> (Accessed: 5 February 2026).
15. Workforce Institute (2023) 'Design standards for older users: High contrast and color differentiation', *Workforce Institute*, 26 September. Available at: <https://workforceinstitute.io/ui-ux-design/design-standards-older-users/> (Accessed: 5 February 2026).